


Faunitaxys

*Revue de Faunistique, Taxonomie et Systématique
morphologique et moléculaire*



Volume 12
Numéro 49

Septembre 2024

ISSN (Print) : 2269 - 6016
ISSN (Online) : 2970 - 4960
Dépôt légal : Septembre 2024

Faunitaxys

*Revue de Faunistique, Taxonomie et Systématique
morphologique et moléculaire*

ZooBank : <https://zoobank.org/79A36B2E-F645-4F9A-AE2B-ED32CE6771CC>

Directeur de la publication, rédacteur, conception graphique et PAO:

Lionel Delaunay

Cette revue ne peut pas être vendue

Elle est distribuée par échange aux institutions (version papier)

et sur simple demande aux particuliers (format PDF)

à l'adresse suivante:

AFCFF (Association française de Cartographie de la Faune et de la Flore)

28, rue Voltaire, F- 42100 Saint Etienne

E-mail: faunitaxys.journal@gmail.com

Elle est disponible librement au téléchargement à partir du site:

<https://faunitaxys.fr/>

La parution de *Faunitaxys* est apériodique

Faunitaxys est indexé dans / *Faunitaxys* is indexed in:

- **Zoological Record**

Articles and nomenclatural novelties are referenced by:

- **ZooBank** (<https://zoobank.org>)

Online Archives:

- **HAL** (<https://hal.archives-ouvertes.fr>)

- **Internet Archive** (<https://archive.org>)

Imprimée sur les presses de SPEED COPIE, 6, rue Tréfilerie, F- 42100 Saint-Etienne

Imprimé le 30 09 2024

Possible estimation of litter size in palaeoburmesebuthid scorpions from Early Cretaceous Burmite (Chelicerata: Scorpiones)

WILSON R. LOURENÇO (1,*) & JÜRGEN VELTEN (2)

(1) Muséum national d'Histoire naturelle, Sorbonne Universités, Institut de Systématique, Evolution, Biodiversité (ISYEB), UMR7205-CNRS, MNHN, UPMC, EPHE, CP 53, 57 rue Cuvier, 75005 Paris, France.

- E-mail : wilson.lourenco@mnhn.fr

- ZooBank : <http://zoobank.org/58448BD6-79D7-46CE-AFDD-91EFF2B7D4EF> – Orcid : <https://orcid.org/0000-0002-2386-363X>

(2) Uhländstr 9 a, 67292 Kirchheimbolanden, Germany.

- ZooBank : <http://zoobank.org/F7A6D8F4-5139-4A78-BC39-AB21BA60FE1C>

* Correspondence.

Keywords:

Scorpiones;
Palaeoburmesebuthidae;
fossil;
litter size;
Burmite.

Abstract. – In previous studies concerning litter size in extant buthoid species of scorpions it was suggested that a number of factors could directly be associated with litter size variation. Litter size should be directly proportional to the size of adult females but inversely proportional to the size of the embryos and pro-juveniles. The large body size of pro-juveniles at birth should be directly associated with a more complete embryonic development, leading to post-embryonic developments comprising smaller numbers of instars but with higher values for the morphometric growths. For fossil scorpions and in particular those from Early Cretaceous nothing could be presumed until now. An original observation of pro-juvenile aggregation for *Betaburmesebuthus* suggests that for micro-buthoid fossil scorpions litter size was extremely reduced but composed of large embryos and pro-juveniles.

Lourenço W. R. & Velten J., 2024. – Possible estimation of litter size in palaeoburmesebuthid scorpions from Early Cretaceous Burmite (Chelicerata: Scorpiones). *Faunitaxys*, 12(49): 1 – 4.

DOI: [https://doi.org/10.57800/faunitaxys-12\(49\)](https://doi.org/10.57800/faunitaxys-12(49))

ZooBank: <https://zoobank.org/5F279A52-DBD0-48CC-8BF4-F32BCF868CD6>

Received: 26/09/2024 – Revised: 27/09/2024 – Accepted: 28/09/2024

Introduction

As already emphasized in previous publications (Lourenço, 2002, 2007, 2018), scorpions are unusual among terrestrial arthropods in several characteristics of their life-history. These particular traits of their reproductive biology lead to approximative interpretations by some early authors (Polis & Sissom, 1990). Nevertheless, some precise studies on scorpion embryology were carried out since the end of the 19th century (Laurie, 1890, 1891, 1896a,b) and these were followed by other publications such as those of Pavlovsky (1924, 1925) and Pflugfelder (1930). Since the middle of the 1970s a new interest on the reproductive biology of scorpions took place, in particular concerning their post-embryonic developments. Very few studies however, were dedicated to litter size in scorpions and only a few isolated publications provided some additional information such as Francke (1981). According to this author, litter size was directly proportional to the size of females and inversely proportional to the size of the pro-juveniles. The size of females and the size of pro-juveniles were found to account for 81% of the variation in litter size among the species studied. Subsequent syntheses proposed by Polis & Sissom (1990) and Lourenço (2002) mainly resumed the known data about litter size in scorpions, without further discussions about the factors that could be responsible for its variability. In a more specific study, Lourenço (2007) focused on micro-buthoid scorpions, in the sensu previously defined by Lourenço (2000). This included in particular the families Buthidae C. L. Koch, 1837 and Microcharmidae Lourenço, 1996. In fact, most of the buthid and microcharmidae species defined as having small-sizes also present

largely reduced litter-sizes generally inferior to 10 (Fig. 1-2) but more frequently with numbers lower than 7 pro-juveniles (Lourenço, 2007).

Results

The specimens investigated are preserved in a very clear round to oval-shaped piece of yellow amber that measures 28 x 20 mm and is about 3 mm thick. Two of the three observed scorpions are totally complete and most of the structures and characters are globally visible. The third scorpion was cut and only part of the mesosoma and metasoma are observable (Fig. 3-6). This is the second known piece of Burmite with several scorpions trapped together. The first recorded case was one concerning *Palaeoburmesebuthus ohlhoffi* Lourenço, 2015. The scorpions were however, very poorly preserved juveniles (Lourenço & Velten, 2019). The observation of a few bothria (not the hairs) allows defining the family and genus as *Palaeoburmesebuthidae* Lourenço, 2015 and *Betaburmesebuthus* Lourenço, 2015. In account of the size and number of teeth of the pectines, the species clearly shows affinities with *Betaburmesebuthus kobberti* Lourenço, 2015.

The two well visible specimens represent one male with large pectines and 21-21 teeth and one female with smaller pectines and 18-17 teeth. Their global size reaches 6.51 mm. Their definition as pro-juveniles (or instar I) is clearly possible by the observation of a number of characters:

- (i) buccal apparatus not opened,
- (ii) chelicerae without the presence of serrulas,
- (iii) aculeus not sharpened and not functional,
- (iv) extremity of tarsi without claws or unguis but with the presence of blunts, cup-shaped tips.

Reviewer: Eric Ythier (BYG Taxa, France) - ZooBank: <http://zoobank.org/06FD0852-A88E-49E5-B8E6-E1494B86C4E1>



This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

Copyright 2024. The Authors. *Faunitaxys* published by Lionel Delaunay on behalf of the AFCFF (Association française de Cartographie de la Faune et de la



Fig. 1. *Microtityus rickyi*, Kjellesvig-Waering, 1966. Female with pro-juveniles.



Fig. 2. *Tityus (Caribetityus) elii* Armas & Marcato Fondur, 1992. Female with pro-juveniles.

The only known specimen (holotype) of *B. kobberti* is a male juvenile with a total length of 10.41 mm. It is possible to suppose that one or two more molts should yet be necessary to reach adult stage. Taking in account that theoretical morphometric growth value (constant) defined by Dyar (1890) and Przibram & Megusar (1912) for the development of arthropods is 1:26, it is possible to expect a final global size for this species equal to 13.11 mm (one extra molt) or 16.53 mm (two extra molts). Moreover, among micro-buthoids, females are normally slightly bigger. Nevertheless, final sizes could yet be more important if morphometric growth values are higher. For some extant micro-buthoids the observed growth values vary from 1:28 to 1:33 (Lourenço, 2002).

For the studied pro-juveniles, which measure 6.51 mm, if their development is based on a morphometric growth value of 1:26, one can expect successive values equal to 8.20, 10.33, 13.02 and 16.40 mm, what presumes a total of four molts to reach adult stage. If the morphometric growth values are higher this could even suggest a total development with a lower number of molts. These hypothetical results appear to be in accordance with those observed for extant micro-buthoid scorpions (Lourenço, 2002, 2007, 2018).

Acknowledgements

We are most grateful to Dr. Ben Smele (Amsterdam, Netherlands) for his full support to this study and to Alexander Beigel (Aachen, Germany) for the preparation of several of the photos which illustrated this article.

References

- Dyar H., 1890. – The number of molts in Lepidopterous larvae. *Psyche*, 5: 420-422.
- Francke O. F., 1981. – Birth behavior and life history of *Diplocentrus spitzeri* Stahnke (Scorpiones: Diplocentridae). *The Southwestern Naturalist*, 25 (4): 517-523.
- Laurie M., 1890. – The embryology of a scorpion (*Euscorpium italicus*). *Quarterly Journal of Microscopic Science*, 31: 105-141.
- Laurie M., 1891. – Some points in the development of *Scorpio fulvipes*. *Quarterly Journal of Microscopic Science*, 32: 587-597.
- Laurie M., 1896a. – Notes on the anatomy of some scorpions, and its bearing on the classification of the order. *Annals and Magazine of Natural History*, ser 6, 17: 185-194.
- Laurie M., 1896b. – Further notes on the anatomy and development of scorpions, and their bearing on the classification of the order. *Annals and Magazine of Natural History*, ser., 6, 18: 121-133.
- Lourenço W. R., 2000. – Panbiogeographie, les familles des scorpions et leur répartition géographique. *Biogeographica*, 76 (1): 21-39.
- Lourenço W. R., 2002. – Reproduction in scorpions, with special reference to parthenogenesis. Pp. 71-85, In: S. Toft & N. Scharff (Eds.), *European Arachnology 2000*. Aarhus University Press, Aarhus.
- Lourenço W. R., 2007. – Litter size in micro-buthoid scorpions (Chelicerata, Scorpiones). *Boletín de la Sociedad Entomológica Aragonesa*, 40: 473-477.




Fig. 3-6. *Betaburmesebuthus kobberti* Lourenço.

3. Pro-juveniles 1 to 3 and syninclusions. 4. Pro-juvenile 3 showing metasoma with not sharp and non-functional aculeus. 5. Male pro-juvenile, ventral aspect showing large pectines. 6. Female pro-juvenile, ventral aspect showing reduced pectines.

- Lourenço W. R., 2018. – Scorpions and life-history strategies: from evolutionary dynamics toward the scorpionism problem. *Journal of Venomous Animals and Toxins including Tropical Diseases*, 24 (19): 1-12.
- Lourenço W. R. & Velten J., 2019. – The first case of aggregation in a Cretaceous amber species of scorpion: *Palaeoburmesebuthus ohlhoffi* Lourenço, 2015 (Scorpiones: Palaeoburmesebuthidae). *Revista Ibérica de Aracnologia*, 35: 49-50.
- Pavlovsky E. N., 1924. – On the morphology of the male genital apparatus in scorpions. *Travaux de la Société Naturaliste de Leningrad*, 53: 76-86.
- Pavlovsky E. N., 1925. – Zur Morphologie des weiblichen Genitalapparats und zur Embryologie der Skorpione. *Annuaire du Musée Zoologique de l'Académie des Sciences de l'URSS.*, Leningrad, 26: 137-205.
- Pflugfelder O., 1930. – Zur Embryologie des Skorpions, *Hormurus australasiae* (F.). *Zeitschrift f. wissenschaftliche Zoologie*, 137: 1-23.
- Polis G. A. & Sissom W. D., 1990. – Life History. Pp. 161-223, In: G. A. Polis (ed.), *The Biology of scorpions*. Stanford Univ. Press, Stanford.
- Przibram H. & Megusar F., 1912. – Wachstumsmessungen an *Sphodromantis bioculata* Burm. 1. Länge und Masse. *Archiv für Entwicklungsmechanik der Organismen* (Wilhelm Roux), 34: 680-741.
- Authors contribution**¹
- Publisher Correspondence.* – WL
- Writing the article.* – WL
- Article proofreading.* – WL
- Bibliographic work.* – WL
- Material study.* – WL & JV
- Author of the figures.* – WL (Fig. 1-2)
- ¹ WL = Wilson R. Lourenço – JV = Jürgen Velten

Résumé

Lourenço W. R. & Velten J., 2024. – Possible estimation de la taille des portées chez les scorpions Palaeoburmesebuthidae de l'ambre birman du Crétacé inférieur (Chelicerata : Scorpiones). *Faunitaxys*, 12(49): 1–4.

Dans des études précédentes concernant la taille des portées chez les buthoid actuels, un certain nombre de facteurs ont pu être reliés à la variation de cette taille. La taille des portées semblerait directement proportionnelle à la taille des femelles adultes mais inversement proportionnelle à la taille des embryons et des pro-juvéniles. La taille plus importante des pro-juvéniles à la naissance semble directement associée à un développement embryonnaire davantage complet, conduisant à un développement post-embryonnaire plus court avec un nombre de stades plus réduit mais avec des taux de croissance plus élevés. Pour les scorpions fossiles et en particulier ceux trouvés dans l'ambre crétacé de Birmanie, aucune connaissance sur ce sujet n'est disponible. Une observation originale concernant une agrégation de pro-juvéniles appartenant au genre *Betaburmesebuthus* semble suggérer que chez les micro-buthoid fossiles la taille des portées était très réduite mais composée de gros embryons et pro-juvéniles.

Mots-clés. – Scorpiones, fossile, ambre Birman, Palaeoburmesebuthidae, taille des portées.

Derniers articles publiés

- Vives E. & Trócoli S., 2024. – Una nueva especie de *Pseudochoromorpha* Breuning, 1936 de Sulawesi, Indonesia (Coleoptera, Cerambycidae: Lamiinae). *Faunitaxys*, 12(22) : 1–3. [https://doi.org/10.57800/faunitaxys-12\(22\)](https://doi.org/10.57800/faunitaxys-12(22))
- Huchet J-B., 2024. – *Nothochodaeus lycaon*, a new species from Palawan Island, Philippines (Coleoptera: Scarabaeoidea: Ochodaeidae). *Faunitaxys*, 12(23) : 1–4. [https://doi.org/10.57800/faunitaxys-12\(23\)](https://doi.org/10.57800/faunitaxys-12(23))
- Lourenço W. R. & Velten J., 2024. – One more particular new species belonging to the genus *Chaerilobuthus* Lourenço & Beigel, 2011 (Scorpiones: Chaerilobuthidae). *Faunitaxys*, 12(24): 1–5. [https://doi.org/10.57800/faunitaxys-12\(24\)](https://doi.org/10.57800/faunitaxys-12(24))
- Háva J., 2024. – A new *Anthrenus* Geoffroy, 1762 species from Iran (Coleoptera: Dermestidae: Megatominae: Anthrenini). *Faunitaxys*, 12(25) : 1–3. [https://doi.org/10.57800/faunitaxys-12\(25\)](https://doi.org/10.57800/faunitaxys-12(25))
- Lourenço W. R. & Velten J., 2024. – An unusual new species of *Betaburmesebuthus* Lourenço, 2015 trapped in Burmite (Scorpiones: Palaeoburmesebuthidae). *Faunitaxys*, 12(26): 1–5. [https://doi.org/10.57800/faunitaxys-12\(26\)](https://doi.org/10.57800/faunitaxys-12(26))
- Gouverneur X. & Chemin G., 2024. – Notes sur le genre *Xenohammus* Schwarzer, 1931 (Coleoptera, Cerambycidae) et description d'une nouvelle espèce du Laos. *Faunitaxys*, 12(27): 1–6. [https://doi.org/10.57800/faunitaxys-12\(27\)](https://doi.org/10.57800/faunitaxys-12(27))
- Opitz W., 2024. – Systematics of the checkered beetle genus *Teneromimus* Gahan (Coleoptera: Cleroidea: Cleridae: Orthopleurinae). *Faunitaxys*, 12(28): 1–16. [https://doi.org/10.57800/faunitaxys-12\(28\)](https://doi.org/10.57800/faunitaxys-12(28))
- Háva J., 2024. – Descriptions of ten new *Trogoderma* Dejean, 1821 species from Australia (Coleoptera: Dermestidae: Megatominae). *Faunitaxys*, 12(29) : 1–7. [https://doi.org/10.57800/faunitaxys-12\(29\)](https://doi.org/10.57800/faunitaxys-12(29))
- Geshev G. A. (Qiao-Zhi Yang), 2024. – A peculiar new genus and species of rhinoceros beetle, *Ceratohyphus luzonicus*, from the Philippines (Coleoptera: Scarabaeidae: Dynastinae). *Faunitaxys*, 12(30) : 1–10. [https://doi.org/10.57800/faunitaxys-12\(30\)](https://doi.org/10.57800/faunitaxys-12(30))
- Gouverneur X., 2024. – Descriptions de deux nouvelles espèces de Cerambycidae (Coleoptera) du Laos. *Faunitaxys*, 12(31): 1–6. [https://doi.org/10.57800/faunitaxys-12\(31\)](https://doi.org/10.57800/faunitaxys-12(31))

- Gouverneur X., 2024. – Description d’une nouvelle espèce du genre *Oculohammus* Breuning & de Jong du Vietnam (Coleoptera, Cerambycidae). *Faunitaxys*, 12(32): 1 – 4. [https://doi.org/10.57800/faunitaxys-12\(32\)](https://doi.org/10.57800/faunitaxys-12(32))
- Porion T. & Audibert C., 2024. – Description d’une nouvelle espèce du genre *Eupholus* Boisduval, 1835 de Papouasie-Nouvelle-Guinée (Coleoptera, Curculionidae, Entiminae). *Faunitaxys*, 12(33): 1 – 4. [https://doi.org/10.57800/faunitaxys-12\(33\)](https://doi.org/10.57800/faunitaxys-12(33))
- Vives E., 2024. – Cerambycoides nuevos o interesantes de Vietnam (Coleoptera, Cerambycidae) (Pars 11). *Faunitaxys*, 12(34) : 1 – 4. [https://doi.org/10.57800/faunitaxys-12\(34\)](https://doi.org/10.57800/faunitaxys-12(34))
- Keith D., Pham P.V. & Sabatinelli G., 2024. – Description d’une remarquable nouvelle espèce du genre *Eotrichia* Medvedev, 1951 (Coleoptera: Scarabaeidae, Melolonthinae, Rhizotrogini). *Faunitaxys*, 12(35): 1 – 4. [https://doi.org/10.57800/faunitaxys-12\(35\)](https://doi.org/10.57800/faunitaxys-12(35))
- Sabatinelli G. & Pham P.V., 2024. – A new species of *Didrepanephorus* Wood-Mason, 1878 from Vietnam with notes on the genus (Coleoptera: Scarabaeidae Rutelinae). *Faunitaxys*, 12(36): 1 – 4. [https://doi.org/10.57800/faunitaxys-12\(36\)](https://doi.org/10.57800/faunitaxys-12(36))
- Laguerre M., 2024. – Description of three new species and a new genus of Lithosiini from Dominican Republic (Lepidoptera: Erebiidae: Arctiinae: Lithosiini). *Faunitaxys*, 12(37): 1 – 9. [https://doi.org/10.57800/faunitaxys-12\(37\)](https://doi.org/10.57800/faunitaxys-12(37))
- Chen Z.-T., 2024. – *Cranopygia guizhouensis* sp. nov., a new earwig species of Pygidicranidae (Insecta: Dermaptera) from China. *Faunitaxys*, 12(38): 1 – 4. [https://doi.org/10.57800/faunitaxys-12\(38\)](https://doi.org/10.57800/faunitaxys-12(38))
- Fei-Yi Xin, Hao He, Xiao-Tian Zhong, Yi-Feng Zhang & Zhi-Hao Qi, 2024. – *Lucanus ludivinae*, a newly recorded stag beetle from Vietnam, with a description of a new subspecies (Coleoptera: Lucanidae: Lucaninae). *Faunitaxys*, 12(39): 1 – 7. [https://doi.org/10.57800/faunitaxys-12\(39\)](https://doi.org/10.57800/faunitaxys-12(39))
- Soldati F., Barnouin T., Van Meer C., Mouttet R., Vincent A., Bonifait S., Barbéris S. & Etienne S., 2024. – First or new records of non-native beetles in France (Coleoptera: Bostrichidae, Carabidae, Dryophthoridae, Monotomidae, Nitidulidae, Tenebrionidae and Zopheridae). *Faunitaxys*, 12(40): 1 – 12. [https://doi.org/10.57800/faunitaxys-12\(40\)](https://doi.org/10.57800/faunitaxys-12(40))
- Laborieux L., 2024. – A new species of *Tityus* Koch, 1836 (Scorpiones: Buthidae), from the Cordillera Oriental (Colombia). *Faunitaxys*, 12(41): 1 – 9. [https://doi.org/10.57800/faunitaxys-12\(41\)](https://doi.org/10.57800/faunitaxys-12(41))
- Delahaye N., Marazzi G., Marazzi V. & Messori P., 2024. – Synonymy and description of a new species in *Xixuthrus* Thomson, 1864 from Indonesia (Coleoptera, Cerambycidae, Prioninae, Macrotomini, Xixuthrina). *Faunitaxys*, 12(42): 1 – 10. [https://doi.org/10.57800/faunitaxys-12\(42\)](https://doi.org/10.57800/faunitaxys-12(42))
- Vitali F., 2024. – Taxonomic notes about some Indo-Australian Lamiini (Coleoptera Cerambycidae). *Faunitaxys*, 12(43): 1 – 11. <https://zoobank.org/4D315C8D-2EB6-4F66-819D-B606FF4ACDA3>
- Fei-Yi Xin, Yi-Feng Zhang & Zhi-Hao Qi, 2024. – *Falcicornis zhongi* Xin & Qi, sp. nov., a new stag beetle species from Vietnam (Coleoptera: Lucanidae: Lucaninae). *Faunitaxys*, 12(44): 1 – 6. [https://doi.org/10.57800/faunitaxys-12\(44\)](https://doi.org/10.57800/faunitaxys-12(44))
- Lanuza-Garay A., Santos-Murgas A. & Cambra R.A., 2024. – Checklist of True Weevils (Coleoptera: Curculionidae) from Rancho Frio, Darien National Park, Panama. *Faunitaxys*, 12(45): 1 – 7. [https://doi.org/10.57800/faunitaxys-12\(45\)](https://doi.org/10.57800/faunitaxys-12(45))
- Lourenço W. R. & Velten J., 2024. – New insights on the diversity of the family Chaerilobuthidae Lourenço & Beigel, 2011 with the description of one new genus and species (Scorpiones). *Faunitaxys*, 12(46): 1 – 6. [https://doi.org/10.57800/faunitaxys-12\(46\)](https://doi.org/10.57800/faunitaxys-12(46))
- Platia G. & Konvička O., 2024. – Description of three new species of click beetles (Coleoptera: Elateridae) with an updated catalogue of Elateridae from the Sultanate of Oman. *Faunitaxys*, 12(47): 1 – 11. [https://doi.org/10.57800/faunitaxys-12\(47\)](https://doi.org/10.57800/faunitaxys-12(47))
- Háva J., 2024. – A new *Mycetophagus* (subgenus *Ilendus* Casey, 1900) species from Iran (Coleoptera: Mycetophagidae). *Faunitaxys*, 12(48) : 1 – 3. [https://doi.org/10.57800/faunitaxys-12\(48\)](https://doi.org/10.57800/faunitaxys-12(48))

Faunitaxys est échangée avec les revues suivantes (« print versions ») :

- Annali del Museo Civico di Storia Naturale G. Doria (Italie)
- Boletín de la Asociación española de Entomología (Espagne)
- Boletín de la Sociedad Andaluza de Entomología (Espagne)
- Bollettino del Museo di Storia Naturale di Venezia (Italie)
- Bulletin de la Société linnéenne de Lyon (France)
- Bulletin of Insectology (Italie)
- Heteropterus Rev. Entomol. (Espagne)
- Israel Journal of Entomology (Israël)
- Klapalekiana (République Tchèque)
- Koleopterologische Rundschau (Allemagne)
- Memorie del Museo Civico di Storia Naturale di Verona (Italie)
- Nova Supplementa Entomologica (Allemagne)
- Proceedings of the Entomological Society of Washington (USA)
- Revue suisse de Zoologie (Suisse)
- Spixiana (Allemagne)
- Stuttgarter Beiträge zur Naturkunde A, Biologie (Allemagne)
- Zoosystematica Rossica (Russie)

Faunitaxys

Volume 12, Numéro 49, Septembre 2024

SOMMAIRE

Possible estimation de la taille des portées chez les scorpions Palaeoburmesebuthidae de l'ambre birman du Crétacé inférieur (Chelicerata : Scorpiones).

Wilson R. Lourenço & Jürgen Velten 1 – 4

CONTENTS

Possible estimation of litter size in palaeoburmesebuthid scorpions from Early Cretaceous Burmite (Chelicerata: Scorpiones).

Wilson R. Lourenço & Jürgen Velten 1 – 4




Illustration de la couverture :

Illustration de la Couverture: *Betaburmesebuthus kobberti* Lourenço; pro-juveniles in the stone - piece of amber (AB).

Crédits:

Wilson R. Lourenço : Fig. 1-2.

Alexander Beigel : Fig. 3-6 & couverture.